

### Acetaminophen Dosage (Tylenol)

Please note: One dose lasts 4-6 hours. Do not give more than 5 doses in 24 hours.

		Infant's Suspension Liquid 160mg/5mL	Children's Suspension Liquid 160mg/5mL	Children's Soft Chewable Tablets 80mg each	Junior Strength Chewable Tablets 160mg each
		Dropperful	Teaspoon	Tablet	Tablet
Weight	Age	(Use Syringe)	(Use dosing cup)		
6-11 lbs	0-3 months	1.25ml			
12-17 lbs	4-11 mos	2.5ml	½ tsp.		
18-23 lbs	12-23 mos	3.75ml	¾ tsp.		
24-35 lbs	2-3 yrs	5ml	1 tsp.	2	
36-47 lbs	4-5 yrs		1½ tsp.	3	
48-59 lbs	6-8 yrs		2 tsp.	4	2
60-71 lbs	9-10 yrs		2 ½ tsp.	5	2 ½
72-95 lbs	11 yrs		3 tsp.	6	3
96 lbs & over	12 yrs				4

**Ibuprofen Dosage** Please Note: One Dose last 6-8 hours. Not to be used under 6 months.

		Infant's Ibuprofen Concentrated Drops 50mg/1.25mL	Children's Ibuprofen Suspension 100mg/5mL	Children's Soft Chewable Tablets 50mg each	Junior Strength Chewable Tablets 100mg each	Junior Strength Ibuprofen Caplets 100mg
		Dropperful	Teaspoon	Tablet	Tablet	Caplet
Weight	Age	(Use dropper)	(Use dosing cup)			
12-17 lbs	6-11 mos	1=(1.25 mL)				
18-23 lbs	12-23 mos	1 ½=(1.875 mL)				
24-35 lbs	2-3 yrs	2=(2.5 mL)	1 tsp.	2		
36-47 lbs	4-5 yrs		1½ tsp.	3		
48-59 lbs	6-8 yrs		2 tsp.	4	2	2
60-71 lbs	9-10 yrs		2 ½ tsp.	5	2 ½	2 ½
72-95 lbs	11 yrs		3 tsp.	6	3	3

- Milliliter is abbreviated as mL; 5mL equals one teaspoon (tsp).
- Don't use household teaspoons, which can vary in size.
- Aspirin should not be used in children to treat fever or pain.
- Always check manufacturer's directions